


Structured Clinical Interview for DSM Dissociative Disorders (SCID-D)

The SCID-D is a semi-structured clinical interview for assessing the presence and severity of dissociative disorders according to DSM criteria. It is used to identify different aspects of dissociation, including amnesia, depersonalization, derealization, confusion and dissociative identity, which facilitates differential diagnosis with other psychological disorders.

Number of questions	276 questions
Estimated duration	90 minutes
Language	English
Original name	Structured Clinical Interview for DSM Dissociative Disorders
Target audience	Adults, adolescents
Authors	Marlene Steinberg (1994)
Category	Dissociative disorders

Administer your questionnaires online

Send your patient a link: they answer from their phone or computer, and you receive the scored results by email.

View the questionnaire on mentaal.org ›

Scoring & interpretation

The interview allows each symptom to be rated according to its frequency and severity. The assessment is carried out using a specific guide: each dissociative domain is analyzed separately and then integrated into an overall diagnostic impression according to DSM criteria.

Usage

This tool isn't available on Mentaal. To use this questionnaire in your practice, you need a valid license. We invite you to send it directly from the publisher's website.

Are you a psychologist?

With Mentaal, get meaningful insights to support your patients.

- Standardized tests
- Automatic scoring
- Results sent by email

Try it for free

See all questionnaires

★★★★★

“Skeptical at first, I now can't do without this tool in my practice.”

 Elodie C. - CBT Psychologist